

GENTLY PEEL BACK LABEL HERE

FOR ANIMAL TREATMENT ONLY
EXELPET® FLEABAN® Insecticidal Spray for Cat and Dogs

2.0 g/L (S)-METHOPRENE, 1.8 g/L PYRETHRINS, 3.6 g/L PIPERONYL BUTOXIDE, 6.0 g/L N-OCTYL BICYCLOHEPTENE DICARBOXIMIDE

Exelpet Fleaban Insecticidal Spray for cats and dogs will prevent fleas (*Ctenocephalides* spp.) from producing viable eggs for up to 6 weeks. It also controls fleas on cats and dogs for up to 7 days and controls adult brown dog ticks (*Rhipicephalus sanguineus*) for up to 7 days.

Does not control Paralysis tick (*Ixodes holocyclus*).

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.


Store below 30°C (Room Temperature) in a dry place.

Exelpet Products
Petcare Place, Wodonga, VIC., 3690.

250mL

APVMA Approval No: 51142/250mL/0507

1098884



FOR ANIMAL TREATMENT ONLY
exelpet®

FLEABAN® INSECTICIDAL SPRAY FOR CATS AND DOGS
2.0 g/L (S)-METHOPRENE, 1.8 g/L PYRETHRINS, 3.6 g/L PIPERONYL BUTOXIDE, 6.0 g/L N-OCTYL BICYCLOHEPTENE DICARBOXIMIDE

- ✓ Controls adult fleas and flea eggs on cats and dogs
 - ✓ Prevents the production of viable eggs
 - ✓ Controls fleas for 7 days
 - ✓ Controls adult brown dog ticks for up to 7 days.
- Does not control Paralysis tick

Suitable for:



KITTEN



ADULT CAT



PUPPY



SMALL DOG

LESS THAN 11kg



MED DOG

WEIGHING BETWEEN 11-23kg



LGE DOG

GREATER THAN 23kg

250mL

FLEA AND TICK CONTROL

CAUTION: Flammable. Keep away from naked flame. For optimum control it is recommended that the pet's environment also be treated with a product containing an Insect Growth Regulator.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

Material Safety Data Sheet: If additional Hazard information is required, refer to the Material Safety Data Sheet. For a copy in Australia phone 1800 633 768

Flea Control Programme

This product alone is not sufficient to ensure complete freedom from fleas as your pet can quickly become reinfested with fleas from other pets or from the surroundings.

To obtain maximum control of fleas remember to:

1. Treat all pets at the same time.
2. Treat your pet's bedding and the environment.
3. Maintain this flea control programme all

year round, particularly throughout the warmer, humid months when fleas are most active.

Guarantee of Satisfaction: If you would like more information on controlling fleas and ticks on your cat or dog, or have any queries of any kind, or you are not completely satisfied with this product, please contact **EXELPET** Customer Enquires. Your feedback is important to us. Phone or Post: **Australia:** Toll Free Phone 1800 806 391; Reply Paid 4 PO Box 153 Wodonga VIC., 3689. Or visit our website: www.exelpet.com.au

MADE IN AUSTRALIA.

®Registered Trademark.

©Exelpet Products.

Exelpet Products
Petcare Place, Wodonga, VIC., 3690.

Dispose of empty container by wrapping with paper and putting in garbage.

Store below 30°C (Room Temperature) in a dry place.

APVMA Approval No: 51142/250mL/0507

FOR ANIMAL TREATMENT ONLY

EXELPET® FLEABAN® Insecticidal Spray for Cats and Dogs

2.0 g/L (S)-METHOPRENE, 1.8 g/L PYRETHRINS, 3.6 g/L PIPERONYL BUTOXIDE, 6.0 g/L N-OCTYL BICYCLOHEPTENE DICARBOXIMIDE
Exelpet Fleaban Insecticidal Spray for cats and dogs will prevent fleas (*Ctenocephalides* spp.) from producing viable eggs for up to 6 weeks. It also controls fleas on cats and dogs for up to 7 days and controls adult brown dog ticks (*Rhipicephalus sanguineus*) for up to 7 days.
Does not control Paralysis tick (*Ixodes holocyclus*).

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

Store below 30°C (Room Temperature) in a dry place.

Exelpet Products
Petcare Place, Wodonga, VIC., 3690.

DIRECTIONS FOR USE:

This product should not be used on new born, sick or convalescing animals.

For animals less than 3 months of age it is suggested that the product is applied using a towel which has been dampened with the spray.

This product should not be sprayed in the eyes or on genitals.

To control fleas for up to 7 days and prevent fleas from producing viable eggs for up to 6 weeks:

*Spray, preferably in open air, at a distance of 15-30cm (6-12 inches) from the animal by squeezing trigger rapidly and continuously.

*Spray from rear of the animal to head, fluffing hair to allow spray to penetrate to the skin. Spray until hair and skin are slightly damp. Alternatively, a cloth that has been dampened with the spray can be used.

*Use weekly or as often as necessary in summer and use at least monthly in Winter. Regular use will prevent fleas from becoming a problem.

This product also controls adult brown dog ticks (for up to 7 days).

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

May cause dribbling and unsteady gait in cats. If these occur wipe off excess product and contact a veterinarian.